

## THE IMPACT OF ALVEOLAR HEMORRHAGE ON LUNG FUNCTION TESTS

Alina V. DOBROTA<sup>1,2✉</sup>, Claudia L. TOMA<sup>2,3</sup>, Ionela N. BELACONI<sup>2,3</sup>, Miron A. BOGDAN<sup>2,3</sup>

<sup>1</sup>Clinical Emergency Hospital of Bucharest, Bucharest, Romania

<sup>2</sup>“Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

<sup>3</sup>National Institute of Pneumology “Marius Nasta”, Bucharest, Romania

Received 03 Nov 2020, Accepted 24 Nov 2020

<https://doi.org/10.31688/ABMU.2020.55.4.09>

### ABSTRACT

**Introduction.** Alveolar hemorrhage syndrome is a severe condition, with a high-risk of death. A timely diagnosis and therapy may be life-saving. The „gold-standard“ method for diagnosing alveolar hemorrhage is bronchoscopy with bronchiolo-alveolar lavage. Pulmonary function tests are useful to appreciate the severity of alveolar hemorrhage.

**The objective of the study** was to measure the impact of alveolar hemorrhage on lung function tests and to detect useful associations between Golde score value and the type of alveolitis with spirometry tests, lung diffusing capacity for carbon monoxide (DLCO) and carbon monoxide transfer coefficient (KCOc), and six minutes walking test (6MWT).

**Material and methods.** We conducted a retrospective study on a sample of 60 patients diagnosed with alveolar hemorrhage syndrome, hospitalized in The National Institute of Pneumology “Marius Nasta” Bucharest, Romania, between June 2010 and February 2019.

**Results.** The results of bronchiolo-alveolar lavage show that macrophage alveolitis is associated with moderate-severe alveolar hemorrhage syndrome, while neutrophilic and lymphocytic alveolitis are associated with mild to moderate alveolar hemorrhage syndrome. Pulmonary function tests (spirometry and complex

### RÉSUMÉ

**L'impact de l'hémorragie alvéolaire sur les tests de la fonction pulmonaire**

**Introduction.** Le syndrome de l'hémorragie alvéolaire est une maladie extrêmement grave, parfois avec un risque élevé de décès. Un diagnostic et un traitement rapides peuvent sauver des vies. La méthode standard d'or pour diagnostiquer l'hémorragie alvéolaire est la bronchoscopie avec lavage bronchiolo-alvéolaire. Les tests de la fonction pulmonaire sont utiles pour apprécier la sévérité de l'hémorragie alvéolaire.

**L'objectif de l'étude** était de mesurer l'impact de l'hémorragie alvéolaire sur les tests de la fonction pulmonaire et de détecter des associations utiles entre la valeur du score de Golde et le type d'alvéolite avec des valeurs de spirométrie, des mesures de la capacité de diffusion pulmonaire du monoxyde de carbone (DLCOc), le coefficient de transfert de monoxyde de carbone (KCOc) et le test de la marche de six minutes (6MWT).

**Matériel et méthodes.** Nous avons mené une étude rétrospective sur un échantillon de 60 patients diagnostiqués avec un syndrome d'hémorragie alvéolaire, hospitalisés à l'Institut National de Pneumologie “Marius Nasta” de Bucarest, Roumanie, entre juin 2010 et février 2019.

✉ Address for correspondence:

Alina DOBROTA  
Clinical Emergency Hospital of Bucharest, Bucharest, Romania  
Address: Calea Floreasca no. 8, 014452, Bucharest, Romania  
Email: [alina.valentina.pirvu@gmail](mailto:alina.valentina.pirvu@gmail)

respiratory tests) showed that only Tiffeneau index, maximal expiratory flow at 50% of vital flow capacity and DLCOc were influenced by the presence of moderate to severe alveolar hemorrhage. The degree of desaturation measured at the end of 6MWT did not correlate with the degree of alveolar hemorrhage severity.

**Conclusions.** An abnormal gas transfer is common in alveolar hemorrhage, an increase of DLCOc value being a sensitive marker for the disease.

**Keywords:** alveolar hemorrhage, Golde score, functional lung tests.

#### Abbreviations list:

BAL - bronchiolo-alveolar lavage

AH - alveolar hemorrhage

FVC - forced vital capacity

FEV- forced expiratory volume

IT - Tiffeneau index

MEF50 - maximal expiratory flow at 50% of vital flow capacity

DLCOc - lung diffusing capacity for carbon monoxide

TLC - total lung capacity

KCOc - carbon monoxide transfer coefficient

6MWT - six minutes walking test

## INTRODUCTION

Alveolar hemorrhage (AH) syndrome is a severe condition, with a high-risk of death<sup>1</sup>. It can appear at any age, often associated with other diseases, such as vasculitis or mixed connective tissue disorders. AH may also be the initial manifestation of an underlying systemic disease<sup>2</sup>. The diagnosis of AH needs a bronchoscopy with bronchiolo-alveolar lavage. Bronchiolo-alveolar lavage (BAL) is used in clinical practice for the diagnosis of certain pulmonary diseases (Table 1)<sup>3</sup>. A normal BAL fluid contains heterogenous populations of macrophages and lymphocytes<sup>4</sup>. When BAL detects an AH syndrome, the severity of the hemorrhage is assessed by Golde score,

**Table 1.** Normal cells of the bronchoalveolar lavage (BAL) liquid<sup>5</sup>

Total cell count	<13x10 <sup>6</sup>
Macrophages	> 84%
Lymphocytes	< 15%
Neutrophils	< 3%
Eosinophils	< 0.5%
Mast cells	< 0.5%
Plasmocyte	0%

**Résultats** Les résultats du lavage broncho-alvéolaire montrent que l'alvéolite macrophagique est associée au syndrome d'hémorragie alvéolaire modérée à sévère, tandis que l'alvéolite neutrophile et lymphocytaire sont associées au syndrome d'hémorragie alvéolaire légère à modérée. Les tests de la fonction pulmonaire (spirométrie et tests respiratoires complexes) ont montré que seuls l'IT, le MEF 50 et le DLCOc étaient influencés par la présence d'une hémorragie alvéolaire modérée ou sévère. Le degré de désaturation mesuré à la fin de 6MWT n'était pas corrélé avec le degré de gravité de l'hémorragie alvéolaire.

**Conclusions.** Un transfert de gaz anormal est une constatation caractéristique dans l'hémorragie alvéolaire, une augmentation de la valeur DLCOc est un marqueur sensible pour la maladie.

**Mots-clés:** hémorragie alvéolaire, score de Golde, tests pulmonaires fonctionnels.

that is based on the hemosiderin content of alveolar macrophages (Table 2).

Hemosiderin-laden macrophages (HLM) in BAL fluid were originally used as diagnostic biomarkers of alveolar hemorrhage<sup>6</sup>, the presence of more than 90% Fe-positive alveolar macrophages (siderophages) confirms a severe AH syndrome<sup>4</sup>. The presence of AH affects the alveolo-capillary gas exchanges, with subsequent changes in spirometry results and complex respiratory tests.

Spirometry is the most common and widely used lung function test, that measures the ability to inhale and exhale air in relation to time. Spirometry is a screening test of general respiratory health and for evaluating subjects whose major complaint is

**Table 2.** Golde Score interpretation.

Normal values	4-25
Mild alveolar hemorrhagic syndrome	25-100
Moderate alveolar hemorrhagic syndrome	100-300
Severe alveolar hemorrhagic syndrome	300-400

dyspnea<sup>7</sup>. The main variables of spirometry are forced vital capacity (FVC) and forced expiratory volume (FEV), maximal expiratory flow at 50% of vital flow capacity (MEF 50) and Tiffeneau index (IT), that represents the FEV1/FVC ratio. Measurements of single breath carbon monoxide transfer factor (DLCOc) and transfer coefficient (KCO) are used widely in the diagnosis and monitoring of respiratory diseases<sup>8</sup>. DLCOc is part of the complex functional tests needed to assess the lung function and aims to measure the diffusion capacity of carbon monoxide through the alveolo-capillary membrane (Table 3).

**Table 3.** Severity and classification of DLCOc reduction<sup>9</sup>

Normal DLCO	>75% of predicted value, up to 140%
Mild	60% to lower limit of normal
Moderate	40% to 60%
Severe	<40%

The six-minutes walking test (6MWT) is a simple test, that has been standardized by the American Thoracic Society<sup>10</sup>, to measure the distance a patient can walk on a flat surface within 6 minutes at his own pace, without speeding at the end of this test. The distance travelled in meters is noted, as well as the appearance of dyspnea, measured with the help of the Borg scale. The 6MWT is a tool for an indirect respiratory function assessment<sup>5</sup>. In order to detect a possible latent respiratory insufficiency, a desaturation at effort by more than 4% compared to the

initial value is considered significant<sup>6</sup>. The predictive value of the 6MWT is limited because of the lack of appropriate standardization<sup>11</sup>.

**THE OBJECTIVE OF THE STUDY** was to measure the impact of AH on lung function tests and to detect useful associations between Golde score value and the type of alveolitis with spirometry tests, lung DLCO and carbon monoxide transfer coefficient (KCOc), and six minutes walking test (6MWT). Other parameter followed in our study was the presence of triad: hemoptysis, dyspnea and pulmonary infiltrates.

**MATERIAL AND METHODS**


We conducted a retrospective study on a sample of 60 patients diagnosed with AH syndrome.

Inclusion criteria: patients with diagnosis of AH syndrome, hospitalized in the National Institute of Pneumology “Marius Nasta“, Bucharest, Romania, between June 2010 and February 2019. All patients included have signed an informed consent that they agree to be included in the research. The study was approved by the ethics committee of the Institute (registration number 849/15.01.2018).

Exclusion criteria: patients with incomplete information in the files; patients who did not sign an informed consent.

**Statistical analysis**

Statistical analysis was performed using SPSS 22.0 software for Windows PC. Frequency and percentage were used to describe nominal/ ordinal variables, average, standard deviation for scale variables (scale type). For the comparative analysis of ordinal variables such as DLCOc, KCOc, alveolitis type,




**Figure 1.** The distribution of patients according to the presence of hemoptysis, anemia and pulmonary infiltrates on chest X-ray.

**Table 4.** Statistical correlation between alveolitis type and AH severity.

		Alveolitis type		
		Neutrophilic n=28	Lymphocytic n=11	Macrophage n=15
Severity	mild	6/21.4%	2/18.2%	1/6.7%
	moderate	16/57.1%	8/72.7%	8/53.3%
	severe	6/21.4%	1/9.1%	6/40%

**Chi-Square Tests**

	Value	diff	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.430 <sup>a</sup>	4	0.351
Likelihood ratio	4.707	4	0.319
Linear-by-linear association	2.178	1	0.140
N of valid cases	54		

**Figure 2.** Correlation between alveolitis type and severity of AH.  
Legend: AH – alveolar hemorrhage.

degree of desaturation depending on the type of alveolitis or the degree of severity, we used the Chi square test. The Kendall's tau and Gamma tests were used to assess the strength of the KCOc impact on the severity. The likelihood ratio was used to assess the chance of having a DLCOc effect on the severity in the absence of a statistically significant chi-square test value. To analyse the differences between the mean values of the spirometric variables, we used the one-way ANOVA test. The calculation of the eta factor allowed a differentiation of the effect according to the degree of severity. A p value  $\leq 0.05$  was considered statistically significant.

**RESULTS**

AH syndrome is characterized clinically and paraclinically by the presence of the triad: hemoptysis,

**Table 5.** Median values of the main spirometric variables.

FVC median value	77.42 $\pm$ 2.96
FEV 1 median value	72.5 $\pm$ 2.65
IT median value	75 $\pm$ 1.47
MEF50 median value	53.9 $\pm$ 3.39

anemia and diffuse bilateral pulmonary infiltrates. Some patients do not present this classical triad, but an AH should be suspected when at least two of three criteria are present<sup>1</sup>. In our study, only one third of patients had hemoptysis and more than a half presented pulmonary infiltrates on chest X-ray (Figure 1).

Table 4 shows that macrophage alveolitis is associated with moderate-severe AH syndrome, while




Figure 3. Statistical correlation between the severity of AH and MEF50 value.

Table 8. Tests of between-subjects effects – dependent variable: IT.

Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared	Observed Power <sup>b</sup>
Corrected Model	654.542 <sup>a</sup>	2	327.271	2.748	.073	.091	.521
Intercept	236517.169	1	236517.169	1985.709	.000	.973	1.000
severity	654.542	2	327.271	2.748	.073	.091	.521
Error	6551.031	55	119.110				
Total	333514.076	58					
Corrected Total	7205.573	57					

neutrophilic and lymphocytic alveolitis are associated with mild to moderate AH syndrome, but without significant differences, due to the small number of patients ( $\chi^2(4)=4.4, p=0.351$ ) (Figure 2).

In the case of MEF50, an increase of the mean level is observed from approximately 48-49 units in case of AH syndrome (mild and moderate degree) to approximately 71 units (in case of severe degree) (Table 5). Application of the ANOVA-one way test indicates a statistically significant difference for  $F(2.54) = 4.01$  and  $p = 0.024$  (Tables 6, 7). Eta-partial square indicates a differentiation effect of about 13% ( $\eta^2 = 0.129$ ), but a reduced test power of only 0.69. The differentiation is stronger in the case of moderate and severe AH syndrome, the mild degree determines a greater variability and a confidence interval of the average difference of MEF50 compared

to the severe degree, with the upper limit close to zero (Figure 3).

In the case of IT, there is an increase in the mean level from about 71-73 units (in the case of mild and moderate AH syndrome) and only about 8 units (in the case of severe degree). Application of the ANOVA-one way test indicates a statistically significant difference for  $F(2.55) = 2.74$  and  $p = 0.05$ . Eta-partial square indicates a differentiation effect of approximately 9% ( $\eta^2 = 0.09$ ) and a reduced test power of only 0.52 (Table 8). The differentiation is stronger in the case of moderate and severe AH syndrome (Figure 4).

In this study, most patients (40%) had slightly lower DLCOc value and more than half of the patients had normal KCOc value (Table 9).




Figure 4. Statistical correlation between the severity of AH and IT values.

Table 9. Complex functional tests main parameters.

Parameter	Value	Number of patients/ percentage
DLCOc	Normal	15 (25%)
	Slightly low	24 (40%)
	Moderately low	14 (23.3%)
	Severely low	7 (11.7%)
KCOc	Normal	33 (55%)
	Low	27 (45%)

Legend. DLCOc –single breath carbon monoxide transfer factor; KCOc- transfer coefficient

Table 10. Correlation between the severity of AH and DLCOc values.

DLCOc value	AH severity		
	Mild N=9	Moderate N=36	Severe N=15
normal	1/11.1	9/25%	1/6.7%
slightly low	4/44.4%	12/33.3%	2/13.3%
moderately low	3/33.3%	5/13.9%	4/26.7%
Severly low	1/11.1%	10/27.8%	8/53.3%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.793 <sup>a</sup>	6	0.134
Likelihood Ratio	10.333	6	0.111
Linear-by-Linear Association	4.218	1	0.040
N of Valid Cases	60		

**Table 11.** Symmetric measures of the statistical tests used

		Value	Asymp. Std. Error <sup>a</sup>	Approx. T <sup>b</sup>	Approx. Sig.
Nominal by Nominal	Phi	0.404			0.134
	Cramer's V	0.286			0.134
	Contingency coefficient	0.375			0.134
Ordinal by Ordinal	Kendall's tau-b	0.236	0.094	2.456	0.014
	Gamma	0.359	0.140	2.456	0.014
	Spearman correlation	0.278	0.109	2.202	0.032 <sup>c</sup>
Interval by Interval	Pearson's R	0.267	0.106	2.113	0.039 <sup>c</sup>
N of valid cases		60			

**Table 12.** Correlation between the severity of AH and KCOc values.

		severity			Total	
		mild	moderate	severe		
KCOc	normal	Count	2	24	7	33
		% within severity	22.2%	66.7%	46.7%	55.0%
decreased	Count	7	12	8	27	
	% within severity	77.8%	33.3%	53.3%	45.0%	
Total	Count	9	36	15	60	
	% within severity	100.0%	100.0%	100.0%	100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.308 <sup>a</sup>	2	.043
Likelihood ratio	6.485	2	.039
Linear-by-linear association	.491	1	.484
N of valid cases	60		

**Table 13.** AH severity and degree of desaturation at 6MWT.

		degree of desaturation			Total
		mild	moderate	severe	
Degree of severity	mild	Count	0	0	2
	moderate	Count	4	4	3
	severe	Count	2	3	1
Total	Count	6	7	6	19

**Chi-Square Tests**

	Value	diff	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.189 <sup>a</sup>	4	.268
Likelihood Ratio	5.526	4	.237
Linear-by-Linear Association	1.886	1	.170
N of Valid Cases	19		

Table 10 shows a higher proportion of the normal and slightly lower level of DLCOc in patients with moderate AH, while in the case of severe AH the proportion of patients with moderate to severe low levels of DLCOc increases. By applying the Gamma test as a measure of prediction, we can say

that the level of DLCOc knowledge improves the prediction for an increased level of AH severity with 35.9% (p=0.014) (Table 11).

From Table 12 we observe an increased proportion of low KCOc levels in patients with mild AH, 77.8% compared to the normal level present in only




Figure 5. Correlation between the severity of AH and the degree of desaturation on 6MWT.

22% of cases. In the group with moderate AH, the inversion of the proportions is observed, 33.3% vs 66.6%. The application of the Chi-square test indicates a statistically significant difference,  $\chi^2 = 6.3$ ,  $p = 0.43$ . A low level of KCOc is not associated with a high degree of severity of AH, while the normal level does not exclude an increased severity.

Out of a total of 60 patients, the 6MWT was performed only in 19 patients. Of these, 6 patients showed significant desaturation (more than 4%) at the end of the test (Table 13). The degree of desaturation did not correlate with the degree of severity of AH ( $\chi^2 (4) = 5.18$ ,  $p = 0.268$ ) (Figure 5).

## DISCUSSION

The typical presentation of AH is a triad of hemoptysis, anemia and pulmonary infiltrates on chest X-ray. In a study of 112 consecutive patients hospitalized for AH, this triad was present only in 34% of patients, although most had severe diseases<sup>12</sup>. In another study, the AH triad was observed in 54% of patients, 67% presented with hemoptysis, 79% with anemia and all patients had new pulmonary infiltrates on chest X-ray at admission<sup>13</sup>. In our study of 60 patients, one third had hemoptysis and more than a half had pulmonary infiltrates on chest X-ray. To confirm the diagnosis, it is necessary to perform bronchoscopy with BAL. Technological advances have made flexible bronchoscopy a very effective tool in the diagnosis of

many respiratory diseases, including conditions associated with AH, and paved the way for many more technological breakthroughs, both diagnostic and therapeutic<sup>14</sup>. Flexible bronchoscopy with BAL was used to document AH in all patients included in the study. A rising red blood cells count in sequential BAL aliquots remaining bloody or becoming bloodier during the BAL procedure<sup>15</sup> is considered diagnosis for the AH syndrome<sup>16</sup>. Macrophages may appear as quickly as within 2 days and may persist for several weeks<sup>17</sup>.

In the present study, BAL fluid cell counts, including the presence of siderophages, the main types of alveolitis (macrophage, neutrophilic and lymphocytic) and Golde score were analyzed and showed that macrophage alveolitis is associated with moderate-severe AH syndrome, while neutrophilic and lymphocytic alveolitis are associated with mild to moderate AH syndrome, but without significant differences, due to the small number of patients. In a study of 94 patients diagnosed with alveolar hemorrhage, BAL fluid analysis showed a predominance of polymorphic neutrophils in specimens from patients with bacterial infections, while specimens from patients with fungal, viral, or *Pneumocystis* infections tended to have increased percentages of lymphocytes<sup>18</sup>. However, two important aspects remain unclear: one would be if the infection occurs at the same time as AH, and the second if the infection is a precipitating factor or contributor to the occurrence of diffuse AH. In patients already diagnosed with Wegener's



granulomatosis or Churg-Straus syndrome, a high iron-positive macrophage count tends to be associated with active disease<sup>19</sup>.

To evaluate the impact of AH on lung function, there were performed several pulmonary function tests. FVC, FEV1, IT, MEF50, KCOc and DLCOc were measured by spirometry and plethysmography. Single-breath DLCOc value was adjusted for hemoglobin. All lung function testing was performed following the standards outlined by the American Thoracic Society/European Respiratory Society.

In a study evaluating the relationship between clinical disease characteristics, pulmonary function tests and high-resolution computed tomography (HRCT) findings in Wegener granulomatosis, 5 years after diagnosis<sup>20</sup>, one-third of patients had abnormal pulmonary function tests findings, consisting in reduced FEV1 value, reduced alveolar diffusion by KCO (transfer coefficient) being most common (24%). In our study, FEV1 value did not correlate with the presence of severe AH, but in the case of MEF50, an increased mean value is observed in moderate and severe AH syndrome.

In the presence of normal lung volumes and spirometry, the DLCO may be the sole abnormal test hinting to a pulmonary vascular disorder like chronic thromboembolic disease, pulmonary hypertension or other causes of pulmonary vascular obliteration<sup>21</sup>. Usually, the DLCOc value increases as the alveoli are distended, because the surface area for gas exchange increases and the alveolar-capillary membrane may become thinner<sup>22</sup>.

A high value of DLCOc has been reported in AH and attributed to increased carbon monoxide uptake by intra-alveolar red blood cells, so DLCOc measurement has been classically considered as a useful diagnostic test in AH. However, a recent study of AH in Goodpasture's syndrome showed that DLCOc was increased in only a quarter of cases, and was reduced in half of them, probably as a result of ventilation/perfusion mismatching<sup>23</sup>. Our study shows a higher proportion of the normal and slightly lower level of DLCOc in patients with moderate AH, while in the case of severe AH the proportion of patients with moderate to severe low levels of DLCOc increases, because of the increased availability of hemoglobin within the alveolar compartment.

Monitoring the DLCOc may be useful in detecting exacerbations of AH in established cases, such as patients with idiopathic pulmonary hemosiderosis or antglomerular basement membrane disease<sup>24</sup>. However, the relationship between iron deposition in the lungs and pulmonary dysfunction remains unclear<sup>25</sup>.

Although gas exchange measurements in interstitial lung disease, including lung pathology

associated with AH, usually are done by measuring the DLCOc, it is unclear whether estimates of gas exchange might be improved by measuring oxygen desaturation during 6MWT. A study of 130 patients diagnosed with sarcoidosis found that DLCOc is a good predictor of the absence of severe gas exchange impairment<sup>26</sup>, the same being valid in the case of AH syndrome. Also, a normal DLCOc value was a good predictor of the absence of severe desaturation during the 6MWT in sarcoidosis<sup>26</sup>. In our study, the degree of desaturation did not correlate with the degree of severity of AH. In another study of 50 patients diagnosed with pulmonary hypertension, a condition that may be associated with AH syndrome, it turned out that DLCOc was not related to exercise capacity, but to oxygen desaturation during exercise. TLC measurements did not provide any additional information in relation to the studied outcomes<sup>27</sup>.

The KCO is a measurement of the constant rate for alveolar uptake of CO during breath-holding in the single breath measurement of DLCOc at full inflation<sup>28</sup>. Some of the most common causes of a lower KCO are diseases with diffuse interstitial impairment, like pulmonary fibrosis, connective tissue diseases, vasculitis. KCOc values higher than the reference value could be present in conditions associated with AH, like pulmonary oedema, congestive heart failure, mitral stenosis, Wegener's granulomatosis, systemic lupus erythematosus, idiopathic haemosiderosis<sup>28</sup>. In all these diseases, the severity of alveolar involvement varies and some normal alveoli survive and contribute to CO uptake<sup>28</sup>.

**THE LIMITS OF THE STUDY.** This study has some limitations because of a small number of patients and also because the results are influenced by the patients' ability to perform respiratory function tests correctly. Further studies are needed on a larger group of patients to obtain more statistically significant results.

## CONCLUSIONS

Our study has found that the severity of the alveolar hemorrhage is correlated only with the values of the IT and MEF 50 in case of moderate or severe AH. In terms of alveolo-capillary diffusion, the low value of DLCOc improves the prediction for an increased severity of AH by 35.9%. Also, a low value of the KCOc is not associated with a high-degree of severity of AH but, at the same time, a normal value of KCOc does not exclude an increased severity of the disease. The degree of desaturation at 6MWT did not correlate with the degree of severity of AH.

### Author Contribution

A.V.D, C.L.T, I.B, A.M.B equally contributed to the design of the study, to patient recruitment, statistical analysis and writing of the manuscript. All authors were responsible for the collection and assembly of the articles/published data, and their inclusion and interpretation in this article. All authors contributed to the critical revision of the manuscript for valuable intellectual content. All authors have read and agreed to the published version of the manuscript.

### Compliance with Ethics Requirements

„The authors declare no conflict of interest regarding this article“

„The authors declare that all the procedures and experiments of this study respect the ethical standards in the Helsinki Declaration of 1975, as revised in 2008(5), as well as the national law. Informed consent was obtained from all the patients included in the study“

„No funding for this study“

### Acknowledgements

None

### REFERENCES

- Lazor R. Alveolar haemorrhage syndromes. *Orphan Lung Dis.* 2011;15-31. doi:10.1183/1025448x.10007310
- Lara AR, Schwarz MI. Diffuse alveolar hemorrhage. *Chest.* 2010;137(5):1164-1171. doi:10.1378/chest.08-2084
- Gabe LM, Baker KM, van Beek EJR, Hunninghake GW, Reinhardt JM, Hoffman EA. Effect of segmental bronchoalveolar lavage on quantitative computed tomography of the lung. *Acad Radiol.* 2011;18(7):876-884.
- Noble B, Du Bois RM, Poulter LW. The distribution of phenotypically distinct macrophage subsets in the lungs of patients with cryptogenic fibrosing alveolitis. *Clin Exp Immunol.* 1989;76(1):41-46.
- Ciolan G, Magheran E, Grigorie V, Badescu C, Teleaga C. Alveolar hemorrhage syndrome - causes and diagnostic methods. *Pneumologia.* 2017;66(4):209-214.
- Fukihara J, Taniguchi H, Ando M, et al. Hemosiderin-laden macrophages are an independent factor correlated with pulmonary vascular resistance in idiopathic pulmonary fibrosis: A case control study. *BMC Pulm Med.* 2017;17(1):1-6.
- Delclaux C, Laveneziana P, Garcia G, Ninot G, Roche N, Morelot-Panzini C. [Pulmonary function testing of dyspnea complaint by the pulmonologist]. *Rev Mal Respir.* 2019;36(4):484-494.
- Robson AG, Innes JA. Short term variability of single breath carbon monoxide transfer factor. *Thorax.* 2001;56(5):358-361.
- Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J.* 2005;26(5):948-968.
- Ruppel GL, Enright PL. Pulmonary function testing. *Respir Care.* 2012;57(1):165-175.
- Lee J, Kim YH, Kang JY, Jegal Y, Park SY. Korean guidelines for diagnosis and management of interstitial lung diseases: Part 3. Idiopathic nonspecific interstitial pneumonia. *Tuberc Respir Dis (Seoul).* 2019;82(4):277-284.
- De Prost N, Parrot A, Cuquemelle E, et al. Diffuse alveolar hemorrhage in immunocompetent patients: Etiologies and prognosis revisited. *Respir Med.* 2012;106(7):1021-1032. doi:10.1016/j.rmed.2012.03.015
- Alexandre AT, Vale A, Gomes T. Diffuse alveolar hemorrhage: How relevant is etiology? *Sarcoidosis Vasc Diffus Lung Dis.* 2019;36(1):47-52.
- Daniels JMA. Flexible bronchoscopy. *JMA.* 2017:1-18.
- Grebski E, Hess T, Hold G, Speich R, Russi E. Diagnostic value of hemosiderin-containing macrophages in bronchoalveolar lavage. *Chest.* 1992;102(6):1794-1799.
- Park MS. Diffuse alveolar hemorrhage. *Tuberc Respir Dis (Seoul).* 2013;74(4):151-162.
- Fishbein GA, Fishbein MC. Lung vasculitis and alveolar hemorrhage: pathology. *Semin Respir Crit Care Med.* 2011;32(3):254-263.
- Kahn FW, Jones JM. Analysis of bronchoalveolar lavage specimens from immunocompromised patients with a protocol applicable in the microbiology laboratory. *J Clin Microbiol.* 1988;26(6):1150-1155.
- Schnabel A, Reuter M, Csernok E, Richter C, Gross WL. Subclinical alveolar bleeding in pulmonary vasculitides: Correlation with indices of disease activity. *Eur Respir J.* 1999;14(1):118-124.
- Koldingsnes W, Jacobsen EA, Sildnes T, Hjalmarsen A, Nossent HC. Pulmonary function and high-resolution CT findings five years after disease onset in patients with Wegener's granulomatosis. *Scand J Rheumatol.* 2005;34(3):220-228.
- Fitting J. Transfer factor for carbon monoxide: a glance behind the scene. *Swiss Med Wkly.* 2004;134:413-418.
- Neder JA, Marillier M, Bernard AC, O'donnell DE. Transfer coefficient of the lung for carbon monoxide and the accessible alveolar volume: clinically useful if used wisely. *Breathe.* 2019;15(1):69-76.
- Lazor R, Bigay-Game' L, Cottin V, et al. Alveolar hemorrhage in anti-basement membrane antibody disease: a series of 28 cases. *Medicine.* 2007; 86: 181-193.
- Greening AP, Hughes JM. Serial estimations of carbon monoxide diffusing capacity in intrapulmonary haemorrhage. *Clin Sci (Lond).* 1981;60(5):507-12.
- Priftis KN, Anthracopoulos MB, Tsakanika C, et al. Quantification of siderophages in bronchoalveolar fluid in transfusional and primary pulmonary hemosiderosis. *Pediatr Pulmonol.* 2006;41(10):972-977.
- Chenivresse C, Boulanger S, Langlois C, Wemeau-Stervinou L, Perez T, Wallaert B. Oxygen desaturation during a 6-minute walk test as a predictor of maximal exercise-induced gas exchange abnormalities in sarcoidosis. *J Thorac Dis.* 2016;8(8):1995-2003.
- Farkhooy A, Bellocchia M, Hedenström H, et al. Lung function in relation to six-minute walk test in pulmonary hypertension. *Eur Clin Respir J.* 2020;7(1).
- Hughes JMB, Pride NB. In defence of the carbon monoxide transfer coefficient KCO (TL/VA). *Eur Respir J.* 2001;17(2):168-174.